

Wreath products with the integers, proper actions and Hilbert space compression

Yves STALDER* and Alain VALETTE

March 19, 2006

Abstract

We prove that the properties of acting metrically properly on some space with walls or some $CAT(0)$ cube complex are closed by taking the wreath product with \mathbb{Z} . We also give a lower bound for the (equivariant) Hilbert space compression of $H \wr \mathbb{Z}$ in terms of the (equivariant) Hilbert space compression of H .

Introduction

A *space with walls*, as defined by Haglund and Paulin [HP98], is a pair (X, \mathcal{W}) where X is a set and \mathcal{W} is a set of partitions of X (called *walls*) into two classes, submitted to the condition that any two points of X are separated by finitely many walls.

The main examples of spaces with walls are given by $CAT(0)$ cube complexes (see [BH99]), i.e. metric polyhedral complexes in which each k -cell is isomorphic to the euclidean cube $[-1/2, 1/2]^k$, and the gluing maps are isometries. Indeed, it is a result of Sageev [Sag95] that hyperplanes in a $CAT(0)$ cube complex endow the set of vertices with a structure of space with walls (see [CN05b] and [Nic04] for more on the relation between spaces with walls and $CAT(0)$ cube complexes).

Our first result is the following:

*Supported by the Swiss FNRS, grant 20-109130

Theorem 0.1 *Suppose that a group H acts metrically properly either on some space with walls, or on some $CAT(0)$ cube complex. Then the wreath product $H \wr \mathbb{Z} := (\bigoplus_{\mathbb{Z}} H) \rtimes \mathbb{Z}$ satisfies the same property.*

Guentner-Kaminker defined the *Hilbert space compression* and the *equivariant Hilbert space compression* for any unbounded metric space (endowed with a group action in the second case) [GK04]. These two numbers are quasi-isometry invariants. Since we will deal with uniformly discrete¹ spaces, the following definitions are equivalent to theirs.

Let (X, d) be a uniformly discrete metric space. We define the *Hilbert space compression* of X as the supremum of the numbers $\alpha \in [0, 1]$ such that there exists a Hilbert space \mathcal{H} , positive constants C_1, C_2 and a map $f : X \rightarrow \mathcal{H}$ with

$$C_1 \cdot d(x, y)^\alpha \leq \|f(x) - f(y)\| \leq C_2 \cdot d(x, y) \quad \forall x, y \in X .$$

It is denoted by $R(X, d)$. If H is a group acting on (X, d) by isometries, the *equivariant Hilbert space compression* of X is the supremum of the numbers $\alpha \in [0, 1]$ such that there exists a Hilbert space \mathcal{H} endowed with an action of H by affine isometries, positive constants C_1, C_2 and a H -equivariant map $f : X \rightarrow \mathcal{H}$ with

$$C_1 \cdot d(x, y)^\alpha \leq \|f(x) - f(y)\| \leq C_2 \cdot d(x, y) \quad \forall x, y \in X .$$

It is denoted by $R_H(X, d)$. One has trivially $R_H(X, d) \leq R(X, d)$.

We may view a group H as a metric space thanks to the word length associated with some (not necessarily finite) generating subset S . We denote then by $R(H, S)$ the Hilbert space compression and by $R_H(H, S)$ the equivariant Hilbert space compression. Note that, in case H is finitely generated, up to quasi-isometry the word metric does not depend on the finite generating set, see [dlH00, Proposition IV.22]. In this case we write $R(H)$ and $R_H(H)$ for the corresponding compressions. We also use these shorter notations in the general case if there is no ambiguity about the generating set. It is a remarkable observation of Gromov (see [dCTV, Proposition 4.4] for a proof) that $R(H) = R_H(H)$ for H finitely generated and amenable.

The first examples of finitely generated groups whose Hilbert space compression is different from 0 and 1 appeared recently in [AGS]: Thompson's

¹That is, there exists a constant $\delta > 0$ such that $d(x, y) \geq \delta$ whenever $x \neq y$.

group F and the wreath product $\mathbb{Z} \wr \mathbb{Z}$ (see the end of Section 3 for more on this). Our next Theorem allows in particular to construct more examples.

Given a generating set S for H , if $\Gamma = H \wr \mathbb{Z}$, we always take $\Sigma = S \cup \{s\}$ as generating set for Γ , where s is the positive generator of \mathbb{Z} .

Theorem 0.2 *Let H be a group, with generating set S and let $\Gamma = H \wr \mathbb{Z}$. The non equivariant and equivariant Hilbert space compressions satisfy:*

$$R(H, S) \geq R(\Gamma, \Sigma) \geq \frac{R(H, S)}{R(H, S) + 1} ;$$

$$R_H(H, S) \geq R_\Gamma(\Gamma, \Sigma) \geq \max \left\{ R_H(H, S) - \frac{1}{2}, \frac{R_H(H, S)}{2R_H(H, S) + 1} \right\} .$$

In order to select the best bound, we mention that one has $t - 1/2 \geq t/(2t+1)$ if and only if $t \geq (1 + \sqrt{5})/4 \cong 0.809\dots$ (for $t \in [0, 1]$). Gromov's remark gives immediately a stronger estimate for the equivariant compression.

Corollary 0.3 *Let H be a finitely generated and amenable group and let $\Gamma = H \wr \mathbb{Z}$. The equivariant Hilbert space compression satisfies:*

$$R_H(H) \geq R_\Gamma(\Gamma) \geq \frac{R_H(H)}{R_H(H) + 1} .$$

The proofs of Theorems 0.1 and 0.2 rest on a similar idea: we express $H \wr \mathbb{Z}$ as an HNN-extension in two different ways, which provide two different actions of $H \wr \mathbb{Z}$ on a tree. In Theorem 0.1 we use the product of these two trees, while in Theorem 0.2 we appeal to the affine actions naturally associated with each of these trees (see section 7.4.1 in [CCJ⁺01]).

1 Preliminaries: wreath products and trees

Let Λ be a group, H a subgroup and $\vartheta : H \rightarrow \Lambda$ an injective homomorphism. The *HNN-extension* with *basis* Λ and *stable letter* t relatively to H and ϑ is defined by $HNN(\Lambda, H, \vartheta) = \langle \Lambda, t \mid t^{-1}ht = \vartheta(h) \forall h \in H \rangle$.

Our definition of graphs and trees are those of [Ser77]. Given an HNN-extension $\Gamma = HNN(\Lambda, H, \vartheta)$, the associated *Bass-Serre tree* is defined by

$$V(T) = \Gamma/\Lambda ; E(T) = \Gamma/H \sqcup \Gamma/\vartheta(H) ; \overline{\gamma H} = \gamma t \vartheta(H) ; \overline{\gamma \vartheta(H)} = \gamma t^{-1} H ;$$

$$(\gamma H)^- = \gamma \Lambda ; (\gamma H)^+ = \gamma t \Lambda ; (\gamma \vartheta(H))^- = \gamma \Lambda ; (\gamma \vartheta(H))^+ = \gamma t^{-1} \Lambda$$

where, given an edge e , its origin is denoted by e^- and its terminal vertex by e^+ . It is a tree [Ser77, Theorem 12]. We turn T to an oriented tree by setting $Ar_+(T) = \Gamma/H$, $(\gamma H)^- = \gamma\Lambda$, $(\gamma H)^+ = \gamma t\Lambda$ and the Γ -action on T preserves this orientation. Moreover we remark that the oriented tree is bi-regular: for each vertex of T the outgoing edges are in bijection with Λ/H and the incoming edges are in bijection with $\Lambda/\vartheta(H)$.

We turn to wreath products. Let G, H be groups. We set

$$\Lambda = H^{(G)} = \bigoplus_{g \in G} H = \{ \lambda : G \rightarrow H \text{ with finite support} \} .$$

The group G acts on Λ by automorphisms: $(g \cdot \lambda)(x) = \lambda(g^{-1}x)$. The *wreath product* $H \wr G$ is the semi-direct product $\Lambda \rtimes G$, with respect to the action above. The group H embeds in $H \wr G$ as the copy of index 1_G . It is easy to see that, given generating sets of G and H , their union generates $H \wr G$.

In case $G = \mathbb{Z}$, one may express $H \wr \mathbb{Z}$ as an HNN-extension in two ways (we denote by s the generator of \mathbb{Z} in $H \wr \mathbb{Z}$ and by t_+, t_- the stable letters of the HNN-extensions)²:

1. Set $\Lambda_+ = \bigoplus_{n \geq 0} H$ and $\vartheta_+ : \Lambda_+ \rightarrow \Lambda_+$ given by $\vartheta_+(\lambda)_0 = 1_H$ and $\vartheta_+(\lambda)_n = \lambda_{n-1}$ for $n \geq 1$. One has $HNN(\Lambda_+, \Lambda_+, \vartheta_+) = H \wr \mathbb{Z}$ and the isomorphism is given by $\lambda \mapsto \lambda$ and $t_+ \mapsto s^{-1}$;
2. Set $\Lambda_- = \bigoplus_{n \leq 0} H$ and $\vartheta_- : \Lambda_- \rightarrow \Lambda_-$ given by $\vartheta_-(\lambda)_0 = 1_H$ and $\vartheta_-(\lambda)_n = \lambda_{n+1}$ for $n \leq -1$. One has $HNN(\Lambda_-, \Lambda_-, \vartheta_-) = H \wr \mathbb{Z}$ and the isomorphism is given by $\lambda \mapsto \lambda$ and $t_- \mapsto s$;

Given a wreath product $H \wr \mathbb{Z}$, we will denote by T_+ , respectively T_- , the Bass-Serre tree associated to the second, respectively third, HNN-extension above. We take as base points (when necessary) the vertices Λ_+ and Λ_- .

We collect now some observations about the $H \wr \mathbb{Z}$ -actions on T_+ and T_- which will be relevant in the next sections. Set $\Gamma = H \wr \mathbb{Z}$ and $\gamma = (\lambda, n) \in \Gamma$. If λ is nontrivial, we set $m = \min\{k \in \mathbb{Z} : \lambda_k \neq 1_H\}$ and $M = \max\{k \in \mathbb{Z} : \lambda_k \neq 1_H\}$.

Lemma 1.1 *If $\lambda = 1$, one has $d_{T_+}(\Lambda_+, \gamma\Lambda_+) = |n| = d_{T_-}(\Lambda_-, \gamma\Lambda_-)$.*

²The most common way is probably the following: set $\vartheta : \Lambda \rightarrow \Lambda$; $\vartheta(\lambda)_n = \lambda_{n-1}$. One has $HNN(\Lambda, \Lambda, \vartheta) = H \wr \mathbb{Z}$ and the isomorphism is given by $\lambda \mapsto \lambda$ and $t \mapsto s^{-1}$. Nevertheless, this expression will be useless in this article.

The proof is obvious.

Lemma 1.2 *If $\lambda \neq 1$, the distances $d_{T_{\pm}}(\Lambda_{\pm}, \gamma\Lambda_{\pm})$ are given by formulas:*

$$\begin{aligned} d_{T_+}(\Lambda_+, \gamma\Lambda_+) &= \begin{cases} |n| & \text{if } n \leq m \text{ or } m \geq 0 \\ n - 2m & \text{if } n > m \text{ and } m < 0 \end{cases} ; \\ d_{T_-}(\Lambda_-, \gamma\Lambda_-) &= \begin{cases} |n| & \text{if } n \geq M \text{ or } M \leq 0 \\ 2M - n & \text{if } n < M \text{ and } M > 0 \end{cases} . \end{aligned}$$

In particular, the inequalities $d_{T_+}(\Lambda_+, \gamma\Lambda_+) \geq -m$, $d_{T_-}(\Lambda_-, \gamma\Lambda_-) \geq M$, $d_{T_+}(\Lambda_+, \gamma\Lambda_+) \geq |n|$ and $d_{T_-}(\Lambda_-, \gamma\Lambda_-) \geq |n|$ hold.

PROOF. We prove the first equality, leaving the second one, which is very similar, to the reader. We remark that $\gamma = \lambda s^n$ and that, for any $k \in \mathbb{Z}$, the stabilizer of the vertex $t_+^k \Lambda_+$ satisfies

$$\text{Stab}(t_+^k \Lambda_+) = t_+^k \Lambda_+ t_+^{-k} = s^{-k} \Lambda_+ s^k = \bigoplus_{i \geq -k} H . \quad (1.3)$$

Suppose first that $m \geq 0$. Then λ stabilizes the vertex Λ_+ , so that we get $d(\Lambda_+, \gamma\Lambda_+) = d(\lambda\Lambda_+, \lambda s^n \Lambda_+) = d(\Lambda_+, s^n \Lambda_+) = |n|$. If $n \leq m$, the vertex $s^n \Lambda_+ = t_+^{-n} \Lambda_+$ is stabilized by λ , so that $d(\Lambda_+, \gamma\Lambda_+) = d(\Lambda_+, s^n \Lambda_+) = |n|$.

It remains to treat the case $n > m$ and $m < 0$. The vertices on the geodesic from $t_+^{-m} \Lambda_+$ to $t_+^{-n} \Lambda_+$ are $t_+^{-m} \Lambda_+, t_+^{-m-1} \Lambda_+, \dots, t_+^{-n} \Lambda_+$. By (1.3), the vertex $t_+^{-m} \Lambda_+$ is stabilized by λ and $t_+^{-m-1} \Lambda_+$ is not. Thus, the geodesic from Λ_+ to $\gamma\Lambda_+$ passes through $t_+^{-m} \Lambda_+$, so that we get

$$d(\Lambda_+, \gamma\Lambda_+) = d(\Lambda_+, t_+^{-m} \Lambda_+) + d(t_+^{-m} \Lambda_+, \gamma\Lambda_+) = -m + n - m = n - 2m .$$

The proof is complete. \square

Let us now state a formula computing the length of an element of $H \wr \mathbb{Z}$, which is a direct consequence of [Par92, Theorem 1.2]. Note that, even if the theorem was stated for finitely generated groups, it also applies in our case.

Proposition 1.4 *Keep the above notations. Let $\gamma = (\lambda, n) \in \Gamma = H \wr \mathbb{Z}$. In case $\lambda = 1$, one has $|\gamma| = |n|$, while in case $\lambda \neq 1$, the length of γ satisfies:*

$$|\gamma| = L_{\mathbb{Z}}(\gamma) + \sum_{i \in \mathbb{Z}} |\lambda_i| ,$$

where $L_{\mathbb{Z}}(\gamma)$ denotes the length of the shortest path starting from 0, ending at n and passing through m and M in the (canonical) Cayley graph of \mathbb{Z} .

The length $L_{\mathbb{Z}}(\gamma)$ appearing in Proposition 1.4 can be estimated as follows:

Proposition 1.5 *Let $\gamma \in \Gamma = H \wr \mathbb{Z}$. The following inequalities hold:*

$$d_{T_{\pm}}(\Lambda_{\pm}, \gamma\Lambda_{\pm}) \leq L_{\mathbb{Z}}(\gamma) \leq d_{T_+}(\Lambda_+, \gamma\Lambda_+) + d_{T_-}(\Lambda_-, \gamma\Lambda_-)$$

PROOF. If $\gamma = (1, n)$, the result is obvious. We suppose now $\gamma = (\lambda, n)$ with $\lambda \neq 1$. The proof is then a distinction of eight cases which are listed in the following tabular:

n	m	M	d_{T_+}	d_{T_-}	$L_{\mathbb{Z}}$	$d_{T_+} + d_{T_-}$
≥ 0	≥ 0	$> n$	n	$2M - n$	$2M - n$	$2M$
≥ 0	≥ 0	$\leq n$	n	n	n	$2n$
≥ 0	< 0	$> n$	$n - 2m$	$2M - n$	$2M - 2m - n$	$2M - 2m$
≥ 0	< 0	$\leq n$	$n - 2m$	n	$n - 2m$	$2n - 2m$
< 0	$< n$	≤ 0	$n - 2m$	$-n$	$n - 2m$	$-2m$
< 0	$\geq n$	≤ 0	$-n$	$-n$	$-n$	$-2n$
< 0	$< n$	> 0	$n - 2m$	$2M - n$	$2M - 2m + n$	$2M - 2m$
< 0	$\geq n$	> 0	$-n$	$2M - n$	$2M - n$	$2M - 2n$

The values of $d_{T_{\pm}}(\Lambda_{\pm}, \gamma\Lambda_{\pm})$ come from Lemma 1.2; those of $L_{\mathbb{Z}}(\gamma)$ are easy to compute. We now observe that the result is true in the eight cases. \square

Combining Propositions 1.4 and 1.5, one obtains immediately:

Corollary 1.6 *Let $\gamma = (\lambda, n) \in \Gamma = H \wr \mathbb{Z}$. The following inequalities hold:*

$$d_{T_{\pm}}(\Lambda_{\pm}, \gamma\Lambda_{\pm}) + \sum_{i \in \mathbb{Z}} |\lambda_i| \leq |\gamma| \leq d_{T_+}(\Lambda_+, \gamma\Lambda_+) + d_{T_-}(\Lambda_-, \gamma\Lambda_-) + \sum_{i \in \mathbb{Z}} |\lambda_i|$$

2 Metrically proper actions

Let us consider a group G , acting by isometries on a metric space X .

Definition 2.1 *The action is metrically proper if, whenever B is a bounded subset of X , the set $\{g \in G : g \cdot B \cap B \neq \emptyset\}$ is finite.*

From now on, we shall write “proper” instead of “metrically proper”. Let us recall that the action is proper if and only if the following property holds, for some $z \in X$:

$$\text{for any } R > 0, \text{ the set } \{g \in G : d(z, g \cdot z) \leq R\} \text{ is finite.} \quad (\text{Prop}_z)$$

Let now $\mathcal{F} = (X_i, b_i)_{i \in I}$ be a family of pointed metric spaces and let $p \geq 1$. We call ℓ^p -product of the family the space

$$\ell^p(\mathcal{F}) = \left\{ x \in \prod_{i \in I} X_i : \sum_{i \in I} d_i(b_i, x_i)^p < +\infty \right\} .$$

It is a metric space with metric $\delta(x, y) = (\sum_{i \in I} d_i(x_i, y_i)^p)^{1/p}$. We set $(b_i)_{i \in I}$ as base point. Consider now the case $(X_i, b_i) = (X, b)$ for all $i \in \mathbb{Z}$. One has:

$$\ell^p(I; X, b) := \ell^p(\mathcal{F}) = \left\{ \phi : I \rightarrow X : \sum_{i \in I} d(b, \phi_i)^p < +\infty \right\} .$$

If a group H acts by isometries on X , the group $H \wr G$ acts by isometries on $\ell^p(G; X, b)$ in the following way:

$$\begin{cases} (\lambda \cdot \phi)_g &= \lambda_g \cdot \phi_g & \text{for } \lambda \in \bigoplus_{g \in G} H ; \\ (g \cdot \phi)_{g'} &= \phi_{g^{-1}g'} & \text{for } g \in G . \end{cases} \quad (2.2)$$

Given G infinite, observe that, even if the action of H is proper, the action of $H \wr G$ on $\ell^p(G; X, b)$ is *not*. Indeed, there is a G -globally fixed point on $\ell^p(G; X, b)$.

Theorem 0.1 will follow from the following statement.

Proposition 2.3 *Let H be a group acting properly on a metric space X , $b \in X$ and $p \geq 1$. Then, the action of $\Gamma = H \wr \mathbb{Z}$ on $T_+ \times T_- \times \ell^p(\mathbb{Z}; X, b)$, where the product is endowed with the ℓ^p metric, is proper.*

PROOF. We are going to prove property (*Prop_z*) for $z = (\Lambda_+, \Lambda_-, (b)_{i \in \mathbb{Z}})$. Thus let $R > 0$ and $A = \{\gamma \in \Gamma : d(z, \gamma \cdot z) \leq R\}$. Take $\gamma = (\lambda, n) \in A$. We have $d_{T_+}(\Lambda_+, \gamma \Lambda_+) \leq R$, $d_{T_-}(\Lambda_-, \gamma \Lambda_-) \leq R$ and $\sum_{i \in \mathbb{Z}} d(b, \lambda_i \cdot b)^p \leq R^p$.

By lemmata 1.1 and 1.2, one has $M \leq R$, $m \geq -R$ (if M and m are defined) and $|n| \leq R$. Set $B = \{h \in H : d(b, h \cdot b) \leq R\}$. It is a finite set since the H -action is proper.

Hence, one has $|n| \leq R$, $\lambda_i = 1_H$ for $|i| > R$ and $\lambda_i \in B$ for $|i| \leq R$. This leaves finitely many choices for γ , and proves thus that A is finite. \square

Remark 2.4 The space $T_+ \times T_- \times \ell^p(\mathbb{Z}; X, b)$ is canonically isometric to the product $\ell^p(\mathcal{F})$ with $I = \{+, -\} \cup \mathbb{Z}$ and \mathcal{F} given by $\mathcal{F}(+) = (T_+, \Lambda_+)$, $\mathcal{F}(-) = (T_-, \Lambda_-)$ and $\mathcal{F}(i) = (X, b)$ for $i \in \mathbb{Z}$.

PROOF OF OF THEOREM 0.1. We recall first that a tree is a CAT(0) cube complex, hence a space with walls.

It is shown in [CMV04, Section 5] that a ℓ^1 -product of spaces with measured walls carries the same structure. Moreover, we remark that, particularizing the construction to spaces with walls, one gets a space with walls. Hence, we get the conclusion for spaces with walls by proposition 2.3.

Given a CAT(0) cube complex Y , we denote by $Y^{(k)}$ the set of k -cells in Y . Take now a family $\mathcal{F} = (X_i, b_i)_{i \in I}$ of CAT(0) cube complexes with $b_i \in X_i^{(0)}$ and set $\mathcal{F}^{(0)} = (X_i^{(0)}, b_i)_{i \in I}$ for $k \in \mathbb{N}$. We are going to construct a subspace X of $\ell^2(\mathcal{F})$ which is a CAT(0) cube complex.

We define first $X^{(0)} = \ell^2(\mathcal{F}^{(0)})$. Since the distance between two distinct vertices is at least 1, one has

$$X^{(0)} = \bigoplus_{i \in I} (X_i^{(0)}, b_i) := \left\{ v \in \prod_{i \in I} X_i^{(0)} : \{i \in I : v_i \neq b_i\} \text{ is finite} \right\}.$$

For $k \geq 1$, we define then the set of k -cells as

$$X^{(k)} = \left\{ c \in \prod_{i \in I} (X_i^{(0)} \cup \dots \cup X_i^{(k)}) : \sum_{i \in I} \dim(c_i) = k \text{ and } \{i \in I : c_i \neq b_i\} \text{ is finite} \right\}.$$

It is clear that every k -cell, as a subset of $\ell^2(\mathcal{F})$, is isometric to $[-1/2, 1/2]^k$. If $c \in X^{(k)}$, the faces of c are the $(k-1)$ -cells c' such that c'_j is a face of c_j for some j and $c'_i = c_i$ for $i \neq j$. The gluing maps are isometric. Finally, the space $\ell^2(\mathcal{F})$ inherits the CAT(0) property, so that X is a CAT(0) cube complex.

Suppose now that H acts on a CAT(0) cube complex Y and take v_0 a vertex of Y . We consider the family \mathcal{F} given by $I = \{+, -\} \cup \mathbb{Z}$, $\mathcal{F}(+) = (T_+, \Lambda_+)$, $\mathcal{F}(-) = (T_-, \Lambda_-)$ and $\mathcal{F}(i) = (Y, v_0)$ for $i \in \mathbb{Z}$. The action of $H \wr \mathbb{Z}$ on $\ell^2(\mathcal{F})$ is proper by proposition 2.3 and the CAT(0) cube complex X constructed as above is an invariant subset, so that it is endowed with a proper action of $H \wr \mathbb{Z}$ too. \square

Remark 2.5 The same techniques can be used, if H acts properly on some Hilbert space \mathcal{H} , to prove that $H \wr \mathbb{Z}$ acts properly on the Hilbert direct sum $\ell^2(E(T_+)) \oplus \ell^2(E(T_-)) \oplus \bigoplus_{i \in \mathbb{Z}} \mathcal{H}$. Hence, we recover the known fact that Haagerup property is preserved by taking wreath products with \mathbb{Z} [CCJ⁺01, Proposition 6.1.1 and Example 6.1.6]. The interest of our technique is that we obtain an *explicit* proper action of $H \wr \mathbb{Z}$, knowing a proper action of H .

Remark 2.6 It is known [CMV04, Theorem 1], that a discrete group satisfies the Haagerup property if and only if it acts properly on some space with measured walls. It follows from Remark 2.5 that whenever H acts properly on a space with measured walls, the same holds for $H \wr \mathbb{Z}$. Again, our techniques give an explicit action, as Theorem 0.1 is also valid for spaces with measured walls.

3 Hilbert space compression: Theorem 0.2

We recall that a map $f : X \rightarrow Y$ between metric spaces is *Lipschitz* if there exists $C > 0$ such that $d_Y(f(x), f(y)) \leq C \cdot d_X(x, y)$ for all $x, y \in X$. Given a Lipschitz map $f : X \rightarrow \mathcal{H}$ (whose range is a Hilbert space), we set³ R_f to be the supremum of the numbers $\alpha \in [0, 1]$ such that there exists $D > 0$ with $D \cdot d_X(x, y)^\alpha \leq \|f(x) - f(y)\|$ for all $x, y \in X$.

Given a generating set S of a group H , we recall our convention to take $\Sigma = S \cup \{s\}$ as generating set for $H \wr \mathbb{Z}$, where s is the positive generator of \mathbb{Z} . In order to simplify notations, we do not mention explicitly S and Σ in this section.

The goal of this section is to prove Theorem 0.2. The key result in this way is the following:

Proposition 3.1 *Let H be a group (with a generating set S) and $\Gamma = H \wr \mathbb{Z}$. Suppose that maps $f : H \rightarrow \mathcal{H}$ and $f_\pm : V(T_\pm) \rightarrow \mathcal{H}_\pm$ are Lipschitz with $R_{f_+} = R_{f_-} > 0$ and $R_f > 0$. Then consider the map*

$$\sigma : \Gamma \rightarrow \mathcal{H}' := \mathcal{H}_+ \oplus \mathcal{H}_- \oplus \bigoplus_{i \in \mathbb{Z}} \mathcal{H} ,$$

where, given $\gamma = (\lambda, n) \in H \wr \mathbb{Z}$, we set $\sigma(\gamma)_\pm = f_\pm(\gamma\Lambda_\pm)$ and $\sigma(\gamma)_i = f(\lambda_i)$ for $i \in \mathbb{Z}$. It satisfies $R_\sigma \geq R_f \cdot R_{f_\pm} / (R_f + R_{f_\pm})$ and $R_\sigma \geq \min\{R_{f_\pm}, R_f - \frac{1}{2}\}$

Moreover, if f is H -equivariant and if f_\pm are Γ -equivariant (with respect to some actions by affine isometries), then there exists a H -action by affine isometries on \mathcal{H}' such that σ is Γ -equivariant.

PROOF. We show first that σ is Lipschitz (the reader could remark that it is trivial if H is a finitely generated group; however, this case is also covered

³It does not coincide with the *asymptotic compression* of f defined in [GK04].

by the proof below). Let us take $C, C_+, C_- > 0$ such that

$$\begin{aligned} \|f(s) - f(t)\| &\leq C \cdot |s^{-1}t| \quad \forall s, t \in H ; \\ \|f_{\pm}(u) - f_{\pm}(v)\| &\leq C_{\pm} \cdot d_{T_{\pm}}(u, v) \quad \forall u, v \in V(T_{\pm}) . \end{aligned}$$

Let $x, y \in \Gamma$. We set $\gamma = x^{-1}y$ and write $x = (\xi, p)$, $y = (\eta, q)$, $\gamma = (\lambda, n)$ in $H \wr \mathbb{Z} = \Lambda \rtimes \mathbb{Z}$ (so that $n = q - p$ and $\lambda_i = \xi_{i-p}^{-1}\eta_i$). One has then

$$\left(\sum_{i \in \mathbb{Z}} \|\sigma(x)_i - \sigma(y)_i\|^2 \right)^{\frac{1}{2}} \leq \sum_{i \in \mathbb{Z}} \|f(\xi_i) - f(\eta_i)\| \leq \sum_{j \in \mathbb{Z}} C \cdot |\lambda_j| \leq C \cdot |\gamma| .$$

Moreover, using Corollary 1.6 for the last step, it comes:

$$\|\sigma(x)_{\pm} - \sigma(y)_{\pm}\| \leq C_{\pm} \cdot d_{T_{\pm}}(x\Lambda_{\pm}, y\Lambda_{\pm}) = C_{\pm} \cdot d_{T_{\pm}}(\Lambda_{\pm}, \gamma\Lambda_{\pm}) \leq C_{\pm} \cdot |\gamma| .$$

Thus, we get finally $\|\sigma(x) - \sigma(y)\| \leq (C_+ + C_- + C) \cdot |x^{-1}y|$, which proves that σ is Lipschitz, as desired.

We now turn to the estimation of R_{σ} . Fix any α, β such that $0 < \alpha < R_f$ and $0 < \beta < R_{f_{\pm}}$. There exists constants $C, C_+, C_- > 0$ such that:

$$\begin{aligned} \|f(s) - f(t)\| &\geq C \cdot |s^{-1}t|^{\alpha} \quad \forall s, t \in H ; \\ \|f_{\pm}(u) - f_{\pm}(v)\| &\geq C_{\pm} \cdot d_{T_{\pm}}(u, v)^{\beta} \quad \forall u, v \in V(T_{\pm}) . \end{aligned}$$

We notice first that σ is injective. More precisely, for any $x, y \in \Gamma$, one has

$$x \neq y \implies \|\sigma(x) - \sigma(y)\| \geq \min\{C, C_+, C_-\} . \quad (3.2)$$

Indeed, we express $x = (\xi, p)$ and $y = (\eta, q)$ as above. If $p \neq q$, we obtain $\|\sigma(x)_{\pm} - \sigma(y)_{\pm}\| \geq C_{\pm} \cdot d_{T_{\pm}}(x\Lambda_{\pm}, y\Lambda_{\pm})^{\beta} \geq C_{\pm}$ and if $\xi_i \neq \eta_i$ for some i , we obtain $\|\sigma(x)_i - \sigma(y)_i\| \geq C \cdot |\xi_i^{-1}\eta_i|^{\alpha} \geq C$.

Let us take x, y and γ as above. According to Corollary 1.6, one (at least) of the following cases occurs. We treat them separately. As the case $x = y$ is trivial, we assume $x \neq y$, that is $|\gamma| \geq 1$, in what follows.

(a) Case $d_{T_+}(\Lambda_+, \gamma\Lambda_+) \geq \frac{1}{3}|\gamma|$: We obtain

$$\begin{aligned} \|\sigma(x) - \sigma(y)\| &\geq \|\sigma(x)_+ - \sigma(y)_+\| \geq C_+ \cdot d_{T_+}(x\Lambda_+, y\Lambda_+)^{\beta} \\ &= C_+ \cdot d_{T_+}(\Lambda_+, \gamma\Lambda_+)^{\beta} \geq \frac{C_+}{3^{\beta}} |x^{-1}y|^{\beta} . \end{aligned}$$

(b) **Case** $d_{T_-}(\Lambda_-, \gamma\Lambda_-) \geq \frac{1}{3}|\gamma|$: We obtain the same way

$$\|\sigma(x) - \sigma(y)\| \geq \|\sigma(x)_- - \sigma(y)_-\| \geq \frac{C_-}{3^\beta} |x^{-1}y|^\beta .$$

(c) **Case** $\sum_{i \in \mathbb{Z}} |\lambda_i| \geq \frac{1}{3}|\gamma|$: We establish two independant estimates.

First, for all $i \in \mathbb{Z}$, one has $\|\sigma(x)_i - \sigma(y)_i\| = \|f(\xi_i) - f(\eta_i)\| \geq C|\xi_i^{-1}\eta_i|^\alpha = C|\lambda_{i+p}|^\alpha$. Hence, using Cauchy-Schwarz inequality for the third step below and $\alpha \leq 1$ for the fourth one, we obtain

$$\begin{aligned} \|\sigma(x) - \sigma(y)\| &\geq \left(\sum_{i \in \mathbb{Z}} \|\sigma(x)_i - \sigma(y)_i\|^2 \right)^{\frac{1}{2}} \geq C \left(\sum_{j \in \mathbb{Z}} |\lambda_j|^{2\alpha} \right)^{\frac{1}{2}} \\ &\geq \frac{C}{\sqrt{M-m+1}} \sum_{j=m}^M |\lambda_j|^\alpha \geq \frac{C}{\sqrt{M-m+1}} \left(\sum_{j=m}^M |\lambda_j| \right)^\alpha \end{aligned}$$

By Proposition 1.4, one has $|\gamma| \geq M - m + 1$, so that we obtain

$$\|\sigma(x) - \sigma(y)\| \geq \frac{C}{\sqrt{|\gamma|}} \left(\frac{1}{3}|\gamma| \right)^\alpha \geq \frac{C}{3^\alpha} |\gamma|^{\alpha - \frac{1}{2}} . \quad (*)$$

This is our first estimate for case (c).

Second, we fix any $\zeta \in]0, 1[$. Then, either there exists $k \in \mathbb{Z}$ such that $|\lambda_k| \geq (\frac{1}{3} \cdot |\gamma|)^\zeta$, or one has $M - m + 1 \geq (\frac{1}{3} \cdot |\gamma|)^{1-\zeta}$. We distinguish the two subcases:

- if there exists $k \in \mathbb{Z}$ such that $|\lambda_k| \geq (\frac{1}{3} \cdot |\gamma|)^\zeta$, we have

$$\begin{aligned} \|\sigma(x) - \sigma(y)\| &\geq \|\sigma(x)_{k-p} - \sigma(y)_{k-p}\| \geq C \cdot |\xi_{k-p}^{-1}\eta_{k-p}|^\alpha \\ &= C \cdot |\lambda_k|^\alpha \geq \frac{C}{3^{\alpha\zeta}} |\gamma|^{\alpha\zeta} ; \end{aligned}$$

- in case $M - m + 1 \geq (\frac{1}{3} \cdot |\gamma|)^{1-\zeta}$, having $L_{\mathbb{Z}}(\gamma) \geq M - m$ by definition, Proposition 1.5 gives

$$d_{T_+}(\Lambda_+, \gamma\Lambda_+) + d_{T_-}(\Lambda_-, \gamma\Lambda_-) \geq L_{\mathbb{Z}}(\gamma) \geq \left(\frac{1}{3} \cdot |\gamma| \right)^{1-\zeta} - 1 .$$

Thus, $\exists s \in \{+, -\}$ such that $d_{T_s}(\Lambda_s, \gamma\Lambda_s) \geq \frac{1}{2}(\frac{1}{3} \cdot |\gamma|)^{1-\zeta} - \frac{1}{2}$. For $|\gamma| \geq 4$, there exists $K > 0$ such that $d_{T_s}(\Lambda_s, \gamma\Lambda_s) \geq K \cdot |\gamma|^{1-\zeta}$, so that $\|\sigma(x) - \sigma(y)\| \geq C_s K^\beta \cdot |x^{-1}y|^{\beta(1-\zeta)}$ as in cases (a)-(b).

Otherwise, for $|\gamma| \leq 3$, equation (3.2) gives

$$\|\sigma(x) - \sigma(y)\| \geq (\min\{C, C_+, C_-\}) \cdot 3^{-\beta(1-\zeta)} \cdot |\gamma|^{\beta(1-\zeta)}.$$

Hence, there exists $C'_\zeta > 0$ with $\|\sigma(x) - \sigma(y)\| \geq C'_\zeta \cdot |x^{-1}y|^{\beta(1-\zeta)}$.

Consequently, setting $m_\zeta = \min\{\alpha\zeta, \beta(1-\zeta)\}$, it comes

$$\|\sigma(x) - \sigma(y)\| \geq \min\left\{\frac{C}{3\alpha\zeta}, C'_\zeta\right\} \cdot |\gamma|^{m_\zeta}. \quad (**_\zeta)$$

The largest value for m_ζ is obtained for $\alpha\zeta = \beta(1-\zeta)$, that is $\zeta = \frac{\beta}{\alpha+\beta}$. It gives $m_\zeta = \frac{\alpha\beta}{\alpha+\beta}$. This is our second estimate for case (c).

As one has $\beta > \alpha\beta/(\alpha+\beta)$, combination of cases (a)-(c) gives

$$\begin{aligned} \|\sigma(x) - \sigma(y)\| &\geq C'' \cdot |x^{-1}y|^{\frac{\alpha\beta}{\alpha+\beta}} && \forall x, y \in \Gamma \\ \|\sigma(x) - \sigma(y)\| &\geq C'' \cdot |x^{-1}y|^{\min\{\beta, \alpha-\frac{1}{2}\}} && \forall x, y \in \Gamma \end{aligned}$$

for some $C'' > 0$. Hence, we get $R_\sigma \geq \alpha\beta/(\alpha+\beta)$ and $R_\sigma \geq \min\{\beta, \alpha-\frac{1}{2}\}$ for all α, β satisfying $0 < \alpha < R_f$ and $0 < \beta < R_{f_\pm}$. This implies immediately $R_\sigma \geq R_f \cdot R_{f_\pm}/(R_f + R_{f_\pm})$ and $R_\sigma \geq \min\{R_{f_\pm}, R_f - \frac{1}{2}\}$.

To conclude the proof of Proposition 3.1, we pass now to the last statement. We thus suppose that f is H -equivariant and f_\pm are Γ -equivariant (with respect to some actions by affine isometries). To establish the Γ -equivariance of σ , we only have to define a Γ -action (by affine isometries) on $\bigoplus_{i \in \mathbb{Z}} \mathcal{H}$ and check the Γ -equivariance with respect to it.

The Γ -action on $\bigoplus_{i \in \mathbb{Z}} \mathcal{H} = \ell^2(\mathbb{Z}, \mathcal{H}, 0)$ is defined by equation (2.2). To check the equivariance, we set $\gamma = (\lambda, n)$ and $g = (\mu, p)$ with $\lambda, \mu \in H^{(\mathbb{Z})}$ and $n, p \in \mathbb{Z}$. We have $(\gamma \cdot \sigma(g))_i = \lambda_i \cdot f(\mu_{i-n})$ and $\sigma(\gamma g)_i = f(\lambda_i \mu_{i-n})$ and we get $(\gamma \cdot \sigma(g))_i = \sigma(\gamma g)_i$ for all i by H -equivariance of f . \square

Theorem 0.2 will be obtained by applying Proposition 3.1 with good embeddings of the trees T_\pm . We explain now how to embed a tree in a Hilbert space with high values of the constant “ R_f ”. First, the following result can be obtained by a straightforward adaptation of [GK04, Proposition 4.2].

Proposition 3.3 *Let $T = (V, E)$ be a tree. Then $R(V) = 1$.*

More precisely, if we denote by E_G the set of geometric (or unoriented) edges of T and if we fix a base vertex v_0 , then for any $\varepsilon \in]0, 1/2[$ we may consider

the map

$$f_\varepsilon : V \longrightarrow \ell^2(E_G) ; x \longmapsto \sum_{k=1}^{d(v_0,x)} k^\varepsilon \delta_{e_k(x)} ,$$

where the $e_k(x)$'s are the consecutive edges on the unique geodesic from v_0 to x and δ_e is the Dirac mass at e . It is a Lipschitz map with $R_{f_\varepsilon} \geq 1/2 + \varepsilon$. We refer to the proof of [GK04, Proposition 4.2] for this fact.

To prove the ‘‘equivariant’’ part of Theorem 0.2, we need some explicit equivariant embeddings into Hilbert spaces. Let $T = (V, E)$ be a tree. We recall from Section 7.4.1 in [CCJ⁺01] how to embed equivariantly T in a Hilbert space. We recall that we denote by $e \mapsto \bar{e}$ the ‘‘orientation-reversing’’ involution on E , and we endow $\ell^2(E)$ with the scalar product:

$$\langle \xi | \eta \rangle = \frac{1}{2} \sum_{e \in E} \xi(e) \overline{\eta(\bar{e})}.$$

Define a map $c : V \times V \rightarrow \ell^2(E) : (x, y) \mapsto c(x, y)$ with

$$c(x, y) = \sum_{e \in (x \rightarrow y)} \delta_e - \delta_{\bar{e}}$$

where δ_e is the Dirac mass at e and the summation is taken over coherently oriented edges in the oriented geodesic from x to y . The map c satisfies, for every $x, y, z \in V$:

$$c(x, y) + c(y, z) = c(x, z); \tag{3.4}$$

$$\|c(x, y)\|^2 = d(x, y). \tag{3.5}$$

Moreover if a group G acts on T , then for every $g \in G$:

$$c(gx, gy) = \pi(g)c(x, y) \tag{3.6}$$

where π is the permutation representation of G on $\ell^2(E)$.

Fix now a base-vertex $v_0 \in V$. Define a map

$$\iota_{T, v_0} : V \rightarrow \ell^2(E) : v \mapsto c(v_0, v)$$

and, for $g \in G$, an affine isometry $\alpha_{v_0}(g)$ of $\ell^2(E)$:

$$\alpha_{v_0}(g)\xi = \pi(g)\xi + c(v_0, gv_0).$$

Using equations (3.4) – (3.6) above, the following lemma is immediate.

- Lemma 3.7**
1. For all $g, h \in G$: $\alpha_{v_0}(gh) = \alpha_{v_0}(g)\alpha_{v_0}(h)$, so that α_{v_0} defines an affine isometric action of G on $\ell^2(E)$;
 2. the map ι_{T, v_0} is G -equivariant with respect to the action α_{v_0} on $\ell^2(E)$;
 3. one has $\|\iota_{T, v_0}(x) - \iota_{T, v_0}(y)\| = \sqrt{d(x, y)}$ for all $x, y \in V$, so that $R_{\iota_{T, v_0}} = 1/2$.

It is an immediate consequence that $R_G(V) \geq 1/2$.

PROOF OF THEOREM 0.2. The inequalities $R(H) \geq R(\Gamma)$ and $R_H(H) \geq R_\Gamma(\Gamma)$ are trivial.

One has $R(V(T_\pm)) = 1$ by Proposition 3.3, so that Proposition 3.1 gives $R(\Gamma) \geq R(V(T_\pm)) \cdot R(H)/(R(V(T_\pm)) + R(H)) = R(H)/(R(H) + 1)$.

Finally, one has $R_\Gamma(V(T_\pm)) \geq 1/2$ by Lemma 3.7, so that we obtain

$$R_\Gamma(\Gamma) \geq \frac{R_\Gamma(V(T_\pm)) \cdot R_H(H)}{R_\Gamma(V(T_\pm)) + R_H(H)} \geq \frac{R_H(H)}{2R_H(H) + 1}$$

$$R_\Gamma(\Gamma) \geq \min \left\{ R_\Gamma(V(T_\pm)), R_H(H) - \frac{1}{2} \right\} = R_H(H) - \frac{1}{2}$$

by Proposition 3.1. □

4 Hilbert space compression: examples

We begin this section with known results about the compression of groups of the form $H \wr \mathbb{Z}$. Let us first state a generalization of [AGS, Theorem 3.9] which gives upper bounds for many of them.

Proposition 4.1 *Let G be a finitely generated group with growth function satisfying $\kappa(n) \asymp n^k$ for some $k > 0$ and let H be a group. We assume the generating set of H chosen such that the word metric is unbounded. Then, the Hilbert space compression of $\Gamma = H \wr G$ satisfies*

$$R(\Gamma, \Sigma) \leq \frac{1 + k/2}{1 + k},$$

where Σ is the union of the generating sets of G and H . In particular, with $G = \mathbb{Z}$, we get $R(H \wr \mathbb{Z}) \leq 3/4$.

The proof is a straightforward adaptation of [AGS, Theorem 3.9].

Remark 4.2 If H is finitely generated, the hypothesis “the word metric is unbounded” means exactly that H is infinite.

Lower bounds, were found by Tessera [Tes, Corollary 14]. In particular:

Proposition 4.3 *Let H be a finitely generated group. If H has polynomial growth, one has $R(H \wr \mathbb{Z}) \geq 2/3$.*

Together, Propositions 4.1 and 4.3 give immediately:

Corollary 4.4 *If H is an infinite group with polynomial growth, then one has $R(H \wr \mathbb{Z}) \in [2/3, 3/4]$.*

In a similar spirit, Proposition 4.1 and our Theorem 0.2 imply immediately:

Corollary 4.5 *Let H be an infinite, finitely generated group.*

- a) *If $R(H) = 1$, then $R(H \wr \mathbb{Z}) \in [1/2, 3/4]$.*
- b) *If $R(H) = R_H(H) = 1/2$, then $R(H \wr \mathbb{Z}) \in [1/3, 1/2]$ and $R_{H \wr \mathbb{Z}}(H \wr \mathbb{Z}) \in [1/4, 1/2]$ (in particular, if $R_{H \wr \mathbb{Z}}(H \wr \mathbb{Z}) < 1/3$, then H is non-amenable).*

□

The interest of part (a) in Corollary 4.5 stems from the fact that numerous groups satisfy $R(H) = 1$: among amenable groups, we mention polycyclic groups and lamplighter groups $F \wr \mathbb{Z}$ with F finite [Tes, Theorem 1]; among (usually) non-amenable groups, we cite hyperbolic groups [BS, Theorem 4.2], groups acting properly co-compactly on finite-dimensional $CAT(0)$ cube complexes [CN05a], co-compact lattices in connected Lie groups, irreducible lattices in higher rank semi-simple Lie groups [Tes, Theorem 2].

Our excuse for isolating (b) in Corollary 4.5 is a remarkable result by Arjantseva, Guba and Sapir [AGS, Theorem 1.8]: for Thompson’s group F , one has $R(F) = R_F(F) = 1/2$.

References

- [AGS] Goulnara N. Arzhantseva, Victor Guba, and Mark Sapir. Metrics on diagram groups and uniform embeddings in a Hilbert space. Preprint arXiv:math.GR/0411605.

- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [BS] Nikolay Brodskiy and Dmitriy Sonkin. Compression of uniform embeddings into Hilbert space. arXiv:math.GR/0509108, preprint, 2005.
- [CCJ⁺01] Pierre-Alain Cherix, Michael Cowling, Paul Jolissaint, Pierre Julg, and Alain Valette. *Groups with the Haagerup property*, volume 197 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2001. Gromov’s a-T-menability.
- [CMV04] Pierre-Alain Cherix, Florian Martin, and Alain Valette. Spaces with measured walls, the Haagerup property and property (T). *Ergodic Theory Dynam. Systems*, 24(6):1895–1908, 2004.
- [CN05a] Sarah Campbell and Graham A. Niblo. Hilbert space compression and exactness of discrete groups. *J. Funct. Anal.*, 222(2):292–305, 2005.
- [CN05b] Indira Chatterji and Graham Niblo. From wall spaces to CAT(0) cube complexes. *Internat. J. Algebra Comput.*, 15(5&6):875–885, 2005.
- [dCTV] Yves de Cornulier, Romain Tessera, and Alain Valette. Isometric group actions on hilbert spaces: growth of cocycles. arXiv:math.GR/0509527, preprint, 2005.
- [dlH00] Pierre de la Harpe. *Topics in geometric group theory*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [GK04] Erik Guentner and Jerome Kaminker. Exactness and uniform embeddability of discrete groups. *J. London Math. Soc. (2)*, 70(3):703–718, 2004.
- [HP98] Frédéric Haglund and Frédéric Paulin. Simplicité de groupes d’automorphismes d’espaces à courbure négative. In *The Epstein*

birthday schrift, volume 1 of *Geom. Topol. Monogr.*, pages 181–248 (electronic). Geom. Topol. Publ., Coventry, 1998.

- [Nic04] Bogdan Nica. Cubulating spaces with walls. *Algebr. Geom. Topol.*, 4:297–309 (electronic), 2004.
- [Par92] Walter Parry. Growth series of some wreath products. *Trans. Amer. Math. Soc.*, 331(2):751–759, 1992.
- [Sag95] Michah Sageev. Ends of group pairs and non-positively curved cube complexes. *Proc. London Math. Soc. (3)*, 71(3):585–617, 1995.
- [Ser77] Jean-Pierre Serre. *Arbres, amalgames, SL_2* . Société Mathématique de France, Paris, 1977. Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass, Astérisque, No. 46.
- [Tes] Romain Tessera. Asymptotic isoperimetry on groups and uniform embeddings into banach spaces. arXiv:math.GR/0603138, preprint, 2006.

Authors address:

Institut de Mathématiques
Université de Neuchâtel
Rue Emile Argand 11
Case postale 158
CH-2009 Neuchâtel
SWITZERLAND

yves.stalder@unine.ch; alain.valette@unine.ch